

CRONO[®] Electrolytes



FICHA TÉCNICA

Electrolito Equilibrado Suplementado con Glucosa, Vitaminas y Minerales

Ingredientes activos (por jeringa):

Glucosa	14.175 mg
Calcio (Ca ⁺⁺)	1.175 mg
Cloruros (Cl ⁻)	4.536 mg
Magnesio (Mg ⁺⁺)	41 mg
Potasio (K ⁺)	1.539 mg
Sodio (Na ⁺)	2.957 mg
Cobalto (Co)	10 mg
Cobre (Cu)	162 mg
Iodo (I)	203 mcg
Hierro (Fe)	243 mg
Manganeso (Mn)	284 mg
Selenio (Se)	24 mcg
Zinc (Zn)	263 mg
Vitamina A	203 UI
Vitamina C	2 mg
Vitamina D ₃	10 UI
Vitamina E	1 mg



Composición: Glucosa, cloruro de sodio, carbonato de calcio, cloruro de potasio, cloruro de magnesio.

Aditivos por kg: Vitaminas – E672 vitamina A 5.000 UI; vitamina C 50 mg; E671 vitamina D₃ 250 UI; 3a700 vitamina E 25 mg. Oligoelementos – E5 Manganeso (quelato de manganeso glicina hidrato) 7.000 mg; E6 Zinc (quelato de zinc glicina hidrato) 6.500 mg; E1 Hierro (quelato de hierro glicina hidrato) 6.000 mg; E4 Cobre (quelato de cobre glicina hidrato) 4.000 mg; E2 Iodo (ioduro de potasio) 5 mg; E8 Selenio (selenito de sodio) 0,6 mg; E3 Cobalto (carbonato de cobalto) 0,25 mg. Aditivos tecnológicos – E321 BHT 0,25 mg/kg.

Propiedades y mecanismo de acción:

- A través del sudor los caballos pierden grandes cantidades de Sodio (Na⁺), Cloro (Cl⁻) y Potasio (K⁺). La pérdida de estos minerales causa fatiga y debilidad muscular, y paradójicamente, disminuye la sensación de sed, bloqueando el mecanismo de rehidratación normal del caballo.
- Los electrolitos son sales minerales que juegan un papel fundamental en el mantenimiento de la presión osmótica, el equilibrio de fluidos corporales, y el funcionamiento óptimo de los músculos y el sistema nervioso.
- Estos minerales median en la transmisión de los impulsos eléctricos para estimular las contracciones musculares.
- Además, un caballo deshidratado tiene menor presión sanguínea, y por tanto un menor flujo sanguíneo en músculos e intestino que puede derivar en cólicos, debilidad y descoordinación muscular, y miositis (acumulación de ácido láctico). Por tanto un aporte adecuado de electrolitos es clave para un óptimo rendimiento de los caballos.
- Por otro lado la deshidratación disminuye la capacidad del caballo para regular su temperatura corporal, causando importantes lesiones en el sistema nervioso.
- Al aportar electrolitos, además de reponer las pérdidas producidas por el sudor, reactivamos el mecanismo de hidratación normal de los caballos, es decir provocamos que los caballos beban y se hidraten adecuadamente.
- La deshidratación es más acusada en ambientes húmedos aunque las temperaturas no sean extremadamente cálidas, por lo que en zonas de alta humedad, con temperaturas por encima de los 12-15°C, el uso de electrolitos es muy recomendable.



Características

Rehidrata y recupera rápidamente al caballo en la competición, entrenamiento intenso, viaje, exposición a altas temperaturas, etc.

Acelera la recuperación muscular tras el ejercicio intenso, lo que permite al caballo volver más rápidamente a su máximo nivel de rendimiento.

Equilibrado: Sodio, Potasio, Magnesio, Calcio y Cloro en proporciones adecuadas.

Suplementado con glucosa, vitaminas A, C, D₃ y E, cobalto, cobre, hierro, iodo, manganeso, selenio y zinc.

Recomendado por veterinarios.

Fácil de administrar.

No contiene sustancias dopantes.

Jeringa oral monodosis.

VetNova

Teléf.: +34 918 440 273 · vetnova@vetnova.net · www.vetnova.net





Electrolito Equilibrado Suplementado con Glucosa, Vitaminas y Minerales

- Más allá de su objetivo fundamental de re-hidratar, CRONO® Electrolytes aporta los siguientes beneficios:
 - Hierro, Cobre y Cobalto estimulan la producción de glóbulos rojos para maximizar el transporte de oxígeno a los músculos, retrasando la aparición de la fatiga y aumentando el rendimiento deportivo.
 - Las vitaminas A, C, D3 y E, Manganeso, Selenio, Yodo y Zinc tienen acción antioxidante: protegen las células y el sistema inmune.
 - La vitamina C es además un eficaz protector articular.
 - La vitamina D es importante para una correcta regulación del Calcio y Fósforo, su absorción intestinal, y para una óptima salud ósea, muscular y tendinosa.
 - Por último, la vitamina E y el Selenio trabajan conjuntamente para reparar el músculo debilitado por el esfuerzo prolongado.

Indicaciones: Ayuda a rehidratar al caballo en situaciones que produzcan una gran pérdida de líquidos y acelera su recuperación muscular tras el ejercicio intenso, lo que le permite volver más rápidamente a su máximo nivel de rendimiento. Situaciones que requieren un aporte regular de electrolitos:

- Ejercicio intenso o prolongado (en cualquier clima o estación).
- Ejercicio ligero en condiciones de elevada humedad y/o temperaturas extremas.
- Exposición prolongada a altas temperaturas (ej: ferias, romerías, etc).
- Caballos en entrenamiento y/o competición en cualquier disciplina.
- Paseos de media-larga distancia.
- Transporte.
- Situaciones de estrés.
- Procesos patológicos que cursen con fiebre elevada y o diarrea.

Especies de destino: Équidos.


Modo de empleo: Inserte la punta de la jeringa en la comisura de la boca del caballo y presione el émbolo asegurándose que ingiere la dosis completa. Si la sudoración ha sido muy intensa, puede ser aconsejable administrar una segunda dosis a las 4-6 horas de la primera. Puede administrarse antes, durante o después de la competición o entrenamiento. Los animales deberán tener acceso permanente al agua.

Advertencias: Este producto no contiene ninguna sustancia prohibida en competición. Guardar el envase bien cerrado, en un lugar fresco, seco, protegido de la luz solar y fuera del alcance de los niños y los animales.

Presentación: Pasta oral en jeringa unidosis (30 ml).

Bibliografía:


- Andrews, F.M., Geiser, D.R., White, S.L., Williamson, L.H., Maykuth, P.L., Greene, E.M., 1995. Haematological and biochemical changes in horses competing in a 3 star horse trial and 3-day-event. *Equine Vet. J. Suppl.* 20, 57-63.
- Andrews, F.M., Ralston, S.L., Williamson, L.H., Maykuth, P.L., White, S.L., Provenza, M., 1995. Weight loss, water loss and cation balance during the endurance test of a three day event. *Equine Vet. J. Suppl.* 18, 294-297.
- Andrews, F.M., Ralston, S.L., Sommardahl, C.S., Maykuth, P.L., Green, E.M., White, S.L., et al., 1994. Weight, water, and cation losses in horses competing in a three-day event. *JAVMA.* 205,721-724.
- Beard, L.A., Hinchliff, K.W., 2002. Effect of NaCl and NaHCO₃ on serum ionised calcium and blood gas status during sprinting. *Equine Vet. J. Suppl.* 34, 519-523.
- Blackmon, D.M., Neathery, M.W., Miller, W.J., Brown, S.R., Carlson, G.P., 1987. Hematology and body fluids in the equine athlete: a review. *Equine Exerc. Physiol.*, San Diego 2, 393-425.
- Carlson, G.P., 2000. Fluid, electrolyte and acid-base balance. In: JJ Kaneko, JW Harvey & ML Bruss (eds), *Clinical Biochemistry of Domestic Animals*. Toronto: Academic Press, pp. 485-516.





Electrolito Equilibrado Suplementado con Glucosa, Vitaminas y Minerales

- Coenen, M., Vervuert, I., 2003. Water and electrolyte turnover in the equine athlete. Uⁿ bers. Tierern7hr. 31, 29–73.
- Coenen, M., 1999. Basics for chloride metabolism and requirement. Proceedings 16th Equine Nutr. Physiol. Symposium, Raleigh, North Carolina, pp. 353–354.
- Coenen, M., Schnermann, J., Vervuert, I., Lindner, A., Sallmann, H.P., 1999a. Effects of oral glucose or electrolyte supplementation after exercise on electrolyte and fluid balance in standardbred horses. Proceedings 16th Equine Nutr. Physiol. Symposium, Raleigh, North Carolina, pp. 304–305.
- Coenen, M., Schuckmann, F., Vervuert, I., von Rennenkampff, F., 1999b. Exercise of pack horses in mountained area. Proceedings 16th Equine Nutr. Physiol. Symposium, Raleigh, North Carolina, pp. 306–307.
- Coenen, M., Meyer, H., Steinbrenner, B., 1995. Effects of NaCl supplementation before exercise on metabolism of water and electrolytes. Equine Vet. J. Suppl. 18, 270–273.
- Coenen, M., 1991a. Chloride Metabolism and Chloride Requirement in Horses. School of Vet. Med. Hannover, Habilitationsschrift. 206 pp.
- Coenen, M., 1991b. Chlorine metabolism in working horses and the improvement of chlorine supply. 12th Proceedings Equine Nutr. Physiol. Symposium, Calgary, Canada, pp. 91–92.
- Coenen, M., Meyer, H., Stadermann, B., 1990. Amount and composition of the GIT content according to type of feed and exercise. Adv. Anim. Physiol. Anim. Nutr. 21, 7–20.
- Dusterdieck, K.F., Schott II, H.C., Eberhart, S.W., Woody, K.A., Coenen, M., 1999. Electrolyte and glycerol supplementation improve water intake by horses performing a simulated 60 km endurance ride. Equine Vet. J. Suppl. 30, 418–424.
- Fedde, M.R., Erickson, H.H., 1998. Increase in blood viscosity in the sprinting horse: can it account for the high pulmonary arterial pressure? Equine Vet. J. 30, 329–334.
- Foreman, J.H., Waldsmith JK., Lalum, R.B., 1999. Environmental stress and three-day eventing: effects of altitude. Equine Vet. J. Suppl. 30, 347–355.
- Foreman, J.H., 1998. The exhausted horse syndrome. Vet. Clin. North Am., Equine Pract. 14, 205–219.
- Foreman, J.H., 1996. Modifications to the 1996 Olympic three-day events to optimize safety under hot and humid conditions. Pferdeheilkunde 12: 397–400.
- Foreman, J.H., 1996. Metabolic causes of equine exercise intolerance. Vet. Clin. North Am., Equine Pract. 12, 537–554.
- Foreman, J.H., 1995. Thermoregulation in the horse exercising under hot and humid conditions. Pferdeheilkunde. 12, 405–408.
- Friend, T.H., 2000. Dehydration, stress, and water consumption of horses during long-distance commercial transport. J. Anim. Sci. 78, 2568–2580.
- Geiser, D.R., Andrews, F.M., Rohrbach, B.W., White, S.L., Maykuth, P.L., Green, E.M., et al., 1995. Blood ionized calcium concentrations in horses before and after the cross-country phase of three-day event competition. American Journal of Veterinary Research 52. 1502–1505.
- Geor, R.J., McCutcheon, L.J., 1998. Hydration effects on physiological strain of horses during exercise—heat stress. J. Appl. Physiol. 84, 2042–2051.
- Geor, R.J., McCutcheon, L.J., 1996. Exercise in the heat: beneficial effects of fluid administration. Journal of Veterinary Internal Medicine 10: 152.
- Geor, R.J., McCutcheon, L.J., Lindinger, M.I., 1996. Adaptations to daily exercise in hot and humid ambient condition-sin trained Thoroughbred horses. Equine Vet. J. Suppl. 22, 63–69.
- Geor, R.J., McCutcheon, L.J., Ecker, G.L., Lindinger, M.I., 1995. Thermal and cardiorespiratory responses of horses to submaximal exercise under hot and humid conditions. Equine Vet. J. Suppl. 20, 125–132.
- Harris PA, Marlin DJ, Mills PC, Roberts CA, Scott CM, Harris RC, et al., (1995). Clinical observations made in non heat acclimated horses performing treadmill exercise in cool (208C/40% RH), hot, dry (308C/40% RH) or hot, humid (308C/80% RH) conditions. Equine Vet. J. Suppl. 20, 78–84.
- Harris PA, Marlin DJ, Scott CM, Harris RC, Mills PC, Michell AR, et al., (1995). Electrolyte and total protein changes in non heat-acclimated horses performing treadmill exercise in cool (208C/40% RH), hot, dry (308C/40% RH) or hot, humid (308C/80% RH) conditions. Equine Vet. J. Suppl. 20, 85–96.
- Harris, P., Snow, D.H., 1992. Plasma potassium and lactate concentrations in thoroughbred horses during exercise of varying intensity. Equine Vet. J. 24, 220–225.
- Hicks JM (1996). Near patient testing: is it here to stay? Journal of Clinical Pathology 49: 191–193.
- Hinchcliff, K.W., 1998. Fluids and electrolytes in athletic horses. Vet. Clin. North Am. Equine Pract., vol. 14:1. Saunders Company, Philadelphia. 225 pp.
- Hinchcliff KW, Kohn CW, Geor R, McCutcheon LJ, Foreman J, Andrews FM, et al. (1995). Acid:base and serum biochemistry changes in horses competing at a modified 1 Star 3-day event. Equine Vet. J. Suppl. 20, 105–110.
- Hyypä, S., Pfsf, A.R., 1998. Fluid, electrolyte, and acid–base responses to exercise in racehorses. Vet. Clin. North






Electrolito Equilibrado Suplementado con Glucosa, Vitaminas y Minerales

Am., Equine Pract. 14 (1), 121–136.

- Hyypä, S., Saastamoinen, M., Pfsf, A.R., 1996. Restoration of water and electrolyte balance in horses after repeated exercise in hot and humid conditions. Equine Vet. J. Suppl. 22, 108–112.
- Jacobs E, Vadasdi E, Sarkozi L and Colman N (1993). Analytical evaluation of I-Stat portable clinical analyser and use by nonlaboratory health-care professionals. Clinical Chemistry 39: 1069–1074.
- Jansson, A., Nyman, S., Morgan, K., Palmgren-Karlsson, C., Lindholm, A., Dahlborn, K., 1995. The effect of ambient temperature and saline loading on changes in plasma and urine electrolytes (Na⁺ and K⁺) following exercise. Equine Vet. J. Suppl. 20, 147–152.
- Jeffcott, L.B., Kohn, C.W., 1999. Contributions of equine exercise physiology research to the success of the 1996 Equestrian Olympic games: a review. Equine Vet. J. Suppl. 30, 347–355.
- Jeffcott, L.B., Clarke, A.F., 1996. Thermoregulatory responses during competitive exercise in the performance horse. Vol II. Equine Vet. J. Suppl., vol. 22. Equine Veterinary Journal, Newmarket, UK. 112 pp.
- Jeffcott, L.B., Clarke, A.F., 1995. Thermoregulatory responses during competitive exercise in the performance horse. Vol I. Equine Vet. J. Suppl., vol. 20. Equine Veterinary Journal, Newmarket, UK. 164 pp.
- Johnson, P.J., 1998. Physiology of body fluids in the horse. Vet. Clin. North Am., Equine Pract. 14 (1), 1–22.
- Jones, J.H., Carlson, G.P., 1995. Estimation of metabolic energy cost and heat production during a 3-day-event. Equine Vet. J. Suppl. 20, 23–30.
- Kingston, J.K., Bayly, W.M., 1998. Effect of exercise on acid–base status of horses. Vet. Clin. North Am., Equine Pract. 14 (1), 61–73.
- Kohn CW, Hinchcliff KW and McKeever K (1999). Effect of ambient temperature and humidity on pulmonary artery temperature of exercising horses. Equine Vet. J. Suppl. 30, 404–411.
- Kohn CW and Hinchcliff KW (1995). Physiological responses to the endurance test of a three-day event during hot and cool weather. Equine Vet. J. Suppl. 20, 31–36.
- Kohn CW, Hinchcliff KW, McCutcheon LJ, Geor R, Foreman J, Allen AK, et al. (1995). Physiological responses of horses competing at a modified 1 star 3-day event. Equine Vet. J. Suppl. 20, 97–104.
- Kohn CW and Brooks CL (1990). Failure of pH to predict ionized calcium percentage in healthy horses. American Journal of Veterinary Research 51: 1206–1210.
- Koterba AM and Carlson GP (1982). Acid–base and electrolyte alterations in horses with exertional rhabdomyolysis. JAVMA. 80, 303–306.
- Kronfeld, D.S., 2001a. Body fluids and exercise: physiological responses: Part 1. J. Equine Vet. Sci. 21, 312–322.
- Kronfeld, D.S., 2001b. Body fluids and exercise: replacement strategies. J. Equine Vet. Sci. 21, 368–375.
- Kronfeld, D.S., 2001c. Body fluids and exercise: influences of nutrition and feeding management. J. Equine Vet. Sci. 21, 417–428.
- Lindinger, M.I., 2004. Acid–base physiology during exercise and in response to training. In: KW Hinchcliff, AJ Kaneps & RJ Geor (eds), Equine Sports Medicine and Surgery: Basic and Clinical Sciences of the Athletic Horse Chapter 39. New York: Elsevier, pp. 586–610.
- Lindinger, M.I., Heigenhauser, G.J., 1991. The roles of ion fluxes in skeletal muscle fatigue. Can. J. Physiol. Pharm. 69, 246–253.
- Marlin, D.J., Scott, C.M., Schroter, R.C., Harris, R.C., Harris, P.A., Roberts, C.A., Mills, P.C., 1999b. Physiological responses of horses to a treadmill simulated speed and endurance test in high heat and humidity before and after humid heat acclimation. Equine Vet. J. 31, 31–42.
- Marlin, D.J., Scott, C.M., Mills, P.C., Loewes, H., Vaarten, J., 1998. Rehydration following exercise: effects of administration of water versus an isotonic oral rehydration solution (ORS). Vet. J. 156, 41–49.
- Marlin, D.J., Scott, C.M., Schroter, R.C., Mills, P.C., Harris, R.C., Harris, P.A., Orme, C.E., Roberts, C.A., Marr, C.M., Dyson, S.J., Barrelet, F., 1996. Physiological responses in nonheat acclimated horses performing treadmill exercise in cool (20°C/40%RH), hot dry (30°C/40%RH) and hot humid (30°C/80%RH) conditions. Equine Vet. J. Suppl. 22, 70–84.
- Marlin, D.J., Harris, P.A., Schroter, R.C., Harris, R.C., Roberts, C.A., Scott, C.M., et al., 1995. Physiological, metabolic and biochemical responses of horses competing in the speed and endurance phase of a CCI***** 3-day event. Equine Vet. J. Suppl. 20: 37–46.
- Masri, M., Freestone, J.F., Wolfsheimer, K.J., Shoemaker, K., 1990. Alterations in plasma volume, plasma constituents, renin activity and aldosterone induced by maximal exercise in the horse. Equine Vet. J. Suppl. 9, 72–77.
- McCutcheon, L.J., Geor, R.J., Ecker, G.L., Lindinger, M.I., 1999. Equine sweating responses to submaximal exercise during 21 days of heat acclimation. J. Appl. Physiol. 87, 1843–1851.
- McCutcheon, L.J., Geor, R.J., 1998. Sweating. Fluid and ion losses and replacement. Vet. Clin. North Am., Equine Pract. 14, 75–95.
- McCutcheon, L.J., Geor, R.J., 1996. Sweat fluid and ion losses in horses during training and competition in cool vs.

VetNova

Teléf.: +34 918 440 273 · vetnova@vetnova.net · www.vetnova.net





Electrolito Equilibrado Suplementado con Glucosa, Vitaminas y Minerales

- hot ambient conditions: implications for ion supplementation. *Equine Vet. J. Suppl.* 22, 54–62.
- McCutcheon, L.J., Geor, R.J., Hare, M.J., Ecker, G.L., Lindiger, M.I., 1995. Sweating rate and sweat composition during exercise and recovery in ambient heat and humidity. *Equine Vet. J. Suppl.* 20, 153–157.
 - McKeever, K.H., 1998a. Blood viscosity and its role in the haemodynamic responses to intense exertion. *Equine Vet. J.* 30, 3.
 - McKeever, K.H., 1998b. Effect of exercise on fluid balance and renal function in horses. *Vet. Clin. North Am., Equine Pract.* 14(1), 23–44.
 - McKeever, K.H., Hinchcliff, K.W., Schmall, L.M., Muir III, W.W., 1991. Renal tubular function in horses during submaximal exercise. *Am. J. Physiol.* 261, R553–R560.
 - Meyer, H., Gomda, Y., Perez-Noriega, H., Heilemann, M., Hipp-Quarton, A., 1990a. Investigation on the postprandial water metabolism in resting and exercised horses. *Adv. Anim. Physiol. Anim. Nutr.* 21, 35–51.
 - Meyer, H., Heilemann, M., Hipp-Quarton, A., Perez-Noriega, H., 1990b. Amount and composition of sweat in ponies. *Adv. Anim. Physiol. Anim. Nutr.* 21, 21–34.
 - Mostert, H.J., Lund, R.J., Guthrie, A.J., Cillers, P.J., 1996. Integrative model for predicting thermal balance in exercising horses. *Equine Vet. J. Suppl.* 22, 7–15.
 - Nyman, S., Jansson, A., Dahlborn, K., Lindholm, A., 1996. Strategies for voluntary rehydration in horses during endurance exercise. *Equine Vet. J. Suppl.* 22, 99–106.
 - Persson, S.G.B., 1967. On blood volume and working capacity in horses. *Acta Vet. Scand. Suppl.* 19 (189 pp.).
 - Rose, R.J., Sampson, D., 1982. Changes in certain metabolic parameters in horses associated with food deprivation and endurance exercise. *Res. Vet. Sci.* 32, 198–202.
 - Rose, R.J., Ilkiw, J.E., 1980. Changes in blood gas, acid–base and metabolic parameters in horses during three-day event competition. *Research in Veterinary Science* 28: 393–395.
 - Rose, R.J., Ilkiw, J.E., Arnold, K.S., Backhouse, J.W., Sampson, D., 1980. Plasma biochemistry in the horse during 3-day event competition. *Equine Vet. J.* 12: 132–136.
 - Schnermann, J., 2000. Effects of oral administration of glucose or electrolyte supplementation after a standardized exercise on thermoregulation and electrolyte metabolism in horses School of Vet. Med. Hannover, thesis, 235 pp.
 - Schott II, H.C., 1998. Oral fluids for equine diarrhoea: an underutilized treatment for a costly disease? *Vet. J.* 155, 119–121.
 - Schott II, H.C., Hinchcliff, K.W., 1998. Treatments affecting fluid and electrolyte status during exercise. *Vet. Clin. North Am., Equine Pract.* 14 (1), 175–204.
 - Schott II, H.C., Hodgson, D.R., Bayly, W., Gollnick, P.D., 1991. Renal response to high intensity exercise. *Equine Exerc. Physiol.* 3, 361–367.
 - Schott II, H.C., Dqsterdieck, K.F., Eberhart, S.W., Woody, K.A., Refsal, K.R., Coenen, M., 1999. Effects of electrolyte and glycerol supplementation on recovery from endurance exercise. *Equine Vet. J. Suppl.* 30, 384–393.
 - Schroter, R.C., Marlin, D.J., 1995. An index of the environmental thermal load imposed on exercising horses and riders by hot weather conditions. *Equine Vet. J. Suppl.* 20, 16–22.
 - Singer, E.R., Saxby, F., French, N.P., 2003. A retrospective case-control study of horse falls in the sport of horse trials and three-day eventing. *Equine Vet. J.* 35, 139–145.
 - Sosa Leon, L.A., Davie, A.J., Hodgson, D.R., Rose, R.J., 1995a. The effects of tonicity, glucose concentration and temperature of an oral rehydration solution on its absorption and elimination. *Equine Vet. J. Suppl.* 20, 140–146.
 - Sosa Leon, L.A., Davie, A.J., Hodgson, D.R., Evans, D.L., Rose, R.J., 1995b. Effect of oral fluid on cardiorespiratory and metabolic responses to prolonged exercise. *Equine Vet. J. Suppl.* 18, 274–278.
 - Taylor, L.E., Ferrante, P.L., Wilson, J.A., Kronfeld, D.S., 1995. Arterial and mixed venous acid–base status and strong ion difference during repeated sprints. *Equine Vet. J. Suppl.* 18, 326–330.
 - Valberg, St.J., Mickelson, J.R., Gallant, E.M., MacLey, J.M., Lentz, L., De La Corte, F., 1999. Exertional rhabdomyolysis in quarter horses and thoroughbreds: one syndrome, multiple aetiologies. *Equine Vet. J. Suppl.* 30, 533–538.
 - Weiss, D.J., Geor, R.J., Burger, K., 1996. Effects of furosemide on hemorheologic alterations induced by incremental treadmill exercise in thoroughbreds. *Am. J. Vet. Res.* 57, 891–895.
 - Williamson, L.H., Andrews, F.M., Maykuth, P.L., White, S.L., Green, E.M., 1996. Biochemical changes in three-day event horses at the beginning, middle and end of Phase C and after Phase D. *Equine Vet. J. Suppl.* 22, 92–98.
 - Williamson, L., White, S., Maykuth, P., Andrews, F., Sommerdahl, C., Green, E., 1995. Comparison between post-exercise cooling methods. *Equine Vet. J. Suppl.* 18, 337–340.

Si le interesa alguno de los artículos listados por favor no dude en solicitarlos a través de los siguientes contactos: vetnova@vetnova.net, 918 440 273 o su Delegad@ Técnico-Comercial VetNova.

VetNova

Teléf.: +34 918 440 273 · vetnova@vetnova.net · www.vetnova.net

